temperature-group

Schauberger

Decades-long observations of Nature, innumerable trials and series of experiments led to the discovery that two fundamentally different **temperature-groups** exist in Nature, namely: [The Energy Evolution - Harnessing Free Energy from Nature, Bio-Technology: Active and Reactive Temperatures]

In Nature both **temperature groups** are active. Group A provides for the progressive build-up of what is suitable for higher development. Group B is responsible for the precipitating out, breaking down and disassembly of everything unfit for such purposes. This has to be removed and as a life-form must be annihilated, or expressed more correctly, reduced to the relatively lowest developmental state, prior to attempting a renewed ascent under entirely different influences. This **temperature-group** also becomes active when any given life-form has fulfilled the purpose of its existence, its duty to multiply itself physically or further develop itself. Withering away through decrepitude, and as an over-ripe product of raw material, it is then made available for the build-up of qualigen with the aid of A-group temperatures (T1 - Schauberger). However, if for any reason B-group temperatures (T2 - Schauberger) become active in the deceased substance, it will be putrefied, combusted or otherwise destroyed. [The Energy Evolution - Harnessing Free Energy from Nature, Bio-Technology: Active and Reactive Temperatures]

In Nature both temperature-groups are active in every healthy organism.

This is necessary because the formative and uplifting levitational essences evolve from the higher temperature group A and the purging dissociative [The Energy Evolution - Harnessing Free Energy from Nature, The Life-Current in Air and Water]

See Also

A-group B-group T1 - Schauberger T2 - Schauberger